

Siberian

Cat breeds: profiles and everyday care



Adult weight (kg)	5.9–11.8
Typical lifespan (years)	11–18+
Coat	Semi-long, dense triple coat with seasonal shedding
Origin	Russia

A practical breed overview. Adult ranges are broad references, not targets, and appearance or breed label cannot predict an individual cat’s health or temperament.

Breed-specific care focus

Comb through the triple coat regularly and check daily during heavy shedding, especially around the belly and hind legs. Provide sturdy climbing routes and play with pauses; adjust for a slow-maturing body and this cat’s mobility.

Health context

This breed is not guaranteed to be safe for people with cat allergy: all cats produce allergens. Ask your vet about this cat’s family health history and preventive care, and an allergy clinician about human symptoms.

A breed can help you prepare, but it is not a diagnosis or a care prescription. Ask your veterinarian to shape an individual plan.

A cat-centred daily rhythm

Food and water

Choose a complete food suited to life stage and health. Measure the daily amount with your veterinary team, divide it into a pattern that suits the cat, and keep fresh water available in a separate, accessible place. Do not infer calories or portions from breed weight ranges.

Litter and shared resources

Keep boxes clean, easy to reach and away from food and noisy traffic. A common starting point is one box per cat plus one extra, placed in separate locations where possible. Scoop at least daily; provide distinct resting, scratching, hiding and feeding areas in multi-cat homes.

Play and enrichment

Offer several short, choice-based play opportunities across the day: wand play, safe searching or food puzzles. Let the cat catch the toy, pause or leave. Add stable vertical space, hiding places and scratching surfaces; rotate toys and supervise strings or small parts.

Medication and handover

Record only medication prescribed for this cat: name, prescribed dose, route, interval, food instructions, dates, and who gave it and when. Check the record before repeating a dose. For a missed dose or reaction, contact the clinic; do not double a dose or change treatment yourself.

Sleep and quiet rest

Cats rest in many short periods across day and night, and the pattern varies with age, health and the home. Provide a warm, quiet, accessible choice of resting places where the cat can retreat undisturbed. A sudden change in sleep, comfort or ability to settle deserves veterinary discussion.

Life stages: adapt the plan

Newborn and very young kitten

From birth, follow a veterinary or experienced rescue plan for warmth, nursing or safe feeding, elimination and individual weight checks. Weakness, feeling cold, difficulty breathing or inability to feed needs prompt veterinary help; an adult schedule must never delay care.

Kitten: birth to 1 year

Support steady growth with a complete kitten diet, frequent age-appropriate meals, safe play and regular veterinary preventive care. Follow the clinic's vaccination, parasite and sterilization advice; activity and feeding needs change quickly as the kitten grows.

Young adult: 1–6 years

Build a stable rhythm around measured complete meals, daily play, clean litter and routine preventive visits. Adjust food and activity to body condition, indoor access, health and the cat's response rather than breed assumptions.

Mature adult: 7–10 years

Keep preventive reviews and note changes in appetite, thirst, weight, grooming, movement or litter habits. Ask the veterinarian whether the check-up schedule or screening should change for this cat.

Senior: over 10 years

Make food, water, litter and favourite resting places easy to reach. Discuss check-up frequency with the veterinarian and record subtle changes in appetite, comfort, sleep, mobility or social behaviour; do not dismiss a new sign as age alone.

Very senior or frail at any age

Prioritise comfort, warmth, low-effort access to essentials and help with grooming or litter when needed. Review pain, night waking and quality of life with the veterinary team; do not apply a younger cat's activity target.

When to contact a veterinarian

Seek emergency veterinary care for open-mouth or difficult breathing, collapse, seizure, major injury or sudden inability to use the back legs. Repeated straining with little or no urine may be a urinary blockage: seek emergency veterinary care immediately; do not wait overnight.

Contact a clinic promptly for not eating, repeated vomiting, marked behaviour change or persistent pain. Kittens, older cats and cats with known illness may need faster assessment.

Care handover log

Task	Time	Caregiver	Notes

Sources and further reading

[TICA — Siberian](#)

[GCCF](#)

[ACAAI](#)

[AAFP/ISFM environmental needs guidelines](#)

[Cornell Feline Health Center: urinary tract disease](#)

[AAHA/AAFP feline life-stage definitions](#)

[Siberian](#)

This guide organizes care observations; it does not diagnose or prescribe. Use the cat's veterinary plan for individual decisions.